

Maximal complexity of finite words

Mira-Cristiana Anisiu* Zoltán Blázsik† Zoltán Kása‡

February 13, 2010

Abstract

The *subword complexity* of a finite word w of length N is a function which associates to each $n \leq N$ the number of all distinct subwords of w having the length n . We define the *maximal complexity* $C(w)$ as the maximum of the subword complexity for $n \in \{1, 2, \dots, N\}$, and the *global maximal complexity* $K(N)$ as the maximum of $C(w)$ for all words w of a fixed length N over a finite alphabet. By $R(N)$ we will denote the set of the values i for which there exists a word of length N having $K(N)$ subwords of length i . $M(N)$ represents the number of words of length N whose maximal complexity is equal to the global maximal complexity.

The values of $K(N)$ and $R(N)$ are obtained; methods to compute $M(N)$ using the de Bruijn graphs and trees are given. An open problem is to find a formula for $M(N)$.

1 Introduction

A *finite word* is a finite sequence of letters over a finite alphabet $\mathcal{A}$, and can be represented as a concatenation of its letters:

$$w = w_1 w_2 \dots w_N \quad \text{with } w_i \in \mathcal{A} \text{ for } 1 \leq i \leq N.$$

The number N is the length of w and is denoted by $|w|$. A word with no letters (i.e. of length 0) is the *empty word*, denoted by ε . We denote by $\mathcal{A}^+$ the set of nonempty words over $\mathcal{A}$, by $\mathcal{A}^* = \mathcal{A}^+ \cup \{\varepsilon\}$ the set of words over $\mathcal{A}$ and by $\mathcal{A}^n$ the set of words of length n over $\mathcal{A}$.

A word u is a *factor* (or *subword*) of w if there exist words $x, y \in \mathcal{A}^*$ such that $w = xuy$. If $x \neq \varepsilon$ and $y \neq \varepsilon$ then u is a *proper factor* (*proper subword*) of w . If $x = \varepsilon$ ($y = \varepsilon$) then u is a *prefix* (*suffix*) of w . Let us

*Tiberiu Popoviciu Institute of Numerical Analysis, Romanian Academy, Cluj-Napoca, E-mail: mira@math.ubbcluj.ro

†Bolyai Institute of Mathematics, University of Szeged, E-mail: blazsik@sol.cc.u-szeged.hu

‡Faculty of Mathematics and Informatics, Babeş-Bolyai University of Cluj-Napoca, E-mail: kasa@cs.ubbcluj.ro

denote by $F(w)$ the set of all nonempty factors of w , and by $F_n(w)$ the set of all factors of w of length n (hence $F_n(w) = F(w) \cap \mathcal{A}^n$).

The *subword complexity* of w counts the number of all distinct factors of a given length occurring in w and is defined as

$$f_w(n) = \text{Card}(F_n(w)) \quad \text{for } 1 \leq n \leq |w|.$$

Clearly $f_w(1) \leq \text{Card}(\mathcal{A})$ and we can consider $f_w(n) = 0$ for $n > |w|$. The subword complexity has been extensively studied in [4], [5] and [6].

The maximal value of the subword complexity $f_w(n)$ for $1 \leq n \leq |w|$ is called the *maximal complexity* of w and is denoted by $C(w)$:

$$C(w) = \max\{f_w(n) \mid n \geq 1\}.$$

The *global maximal complexity* in $\mathcal{A}^N$ is equal to

$$K(N) = \max\{C(w) \mid w \in \mathcal{A}^N\}.$$

We shall denote by $R(N)$ the set of values i for which there exists a word $w \in \mathcal{A}^N$ such that $f_w(i) = K(N)$:

$$R(N) = \{i \in \{1, 2, \dots, N\} \mid \exists w \in \mathcal{A}^N : f_w(i) = K(N)\}.$$

The number of words in $\mathcal{A}^N$ with the maximal complexity equal to the global maximal complexity will be denoted by $M(N)$:

$$M(N) = \text{Card}(\{w \in \mathcal{A}^N : C(w) = K(N)\}).$$

Remark 1. If $\text{Card}(\mathcal{A}) = q$, for $q = 1$ the only word of length N is $w_0 = \underbrace{0 \dots 0}_N$ for which $f_{w_0}(i) = 1$, $i \in \{1, 2, \dots, N\}$, hence $C(w) = 1 =$

$K(N)$, $R(N) = \{1, 2, \dots, N\}$ and $M(N) = 1$. For $q \geq 2$, but $N \leq q$, for each word w_1 which contains N distinct elements of $\mathcal{A}$ we have $C(w_1) = f_{w_1}(1) = N = K(N)$, $R(N) = \{1\}$ and $M(N) = P_q^N$ (permutations of N elements taken from q).

Some values for $K(N)$, $R(N)$ and $M(N)$ in the case of an alphabet of 2 letters are given in Table 1. In the case $N = 3$ the following six words have maximal complexity: 001, 010, 011, 100, 101, 110. For each of them $f_w(1) = 2$, $f_w(2) = 2$, $f_w(3) = 1$, so $K(3) = 2$, $R(3) = \{1, 2\}$ and $M(3) = 6$.

2 Global maximal subword complexity of finite words

In this section we shall compute the values of the global maximal complexity $K(N)$, as well as those of $R(N)$, proving that they are in agreement with the values in Table 1. Some special cases being solved in Remark 1, in what follows we shall consider alphabets with $\text{Card}(\mathcal{A}) = q \geq 2$ and words of length $N > q$.

We shall use the following result.

N	$K(N)$	$R(N)$	$M(N)$
1	1	1	2
2	2	1	2
3	2	1, 2	6
4	3	2	8
5	4	2	4
6	4	2, 3	36
7	5	3	42
8	6	3	48
9	7	3	40
10	8	3	16
11	8	3, 4	558
12	9	4	718
13	10	4	854
14	11	4	920
15	12	4	956
16	13	4	960
17	14	4	912
18	15	4	704
19	16	4	256
20	16	4, 5	79006

Table 1

Lemma 1. [7] *For each $k \in N^*$, the shortest word containing all the q^k words of length k has $q^k + k - 1$ letters (hence in this word each of the q^k words of length k appears only once).*

An algorithm for obtaining such a word for $\mathcal{A} = \{e_1, e_2, \dots, e_q\}$ is the following [7]:

- i.* Each of the first $k - 1$ symbols is equal to e_1 .
- ii.* If the sequence $a_1 a_2 \dots a_k \dots a_{m-k+1} \dots a_{m-1}$ (with $a_1 = \dots = a_{k-1} = e_1$, $m \geq k$ and the a 's representing the e 's in a certain order) has been obtained, the symbol a_m to be added is the e_i with the greatest subscript possible such that $a_{m-k+1} \dots a_{m-1} a_m$ does not duplicate a previously occurring section of k symbols in the above sequence.
- iii.* Rule *ii* is first applied for $m = k$ (in which case $a_m = a_k = e_q$) and then applied repeatedly until a further application is impossible.

Proposition 1. *If $\text{Card}(\mathcal{A}) = q$ and $q^k + k \leq N \leq q^{k+1} + k$ then $K(N) = N - k$.*

Proof. Let us consider at first the case $N = q^{k+1} + k$, $k \geq 1$.

From Lemma 1 we obtain the existence of a word W of length $q^{k+1} + k$ which contains all the q^{k+1} words of length $k+1$, hence $f_W(k+1) = q^{k+1}$. It is obvious that $f_W(l) = q^l < f_W(k+1)$ for $l \in \{1, 2, \dots, k\}$ and $f_W(k+1+j) = q^{k+1} - j < f_W(k+1)$ for $j \in \{1, 2, \dots, q^{k+1} - 1\}$. Any other word of length $q^{k+1} + k$ will have the maximal complexity less than or equal to $C(W) = f_W(k+1)$, hence we have $K(N) = q^{k+1} = N - k$.

For $k \geq 1$ we consider now the values of N of the form $N = q^{k+1} + k - r$ with $r \in \{1, 2, \dots, q^{k+1} - q^k\}$, hence $q^k + k \leq N < q^{k+1} + k$. If from the word W of length $q^{k+1} + k$ considered above we delete the last r letters, we obtain a word W_N of length $N = q^{k+1} + k - r$ with $r \in \{1, 2, \dots, q^{k+1} - q^k\}$. This word will have $f_{W_N}(k+1) = q^{k+1} - r$ and this value will be its maximal complexity. Indeed, it is obvious that $f_{W_N}(k+1+j) = f_{W_N}(k+1) - j < f_{W_N}(k+1)$ for $j \in \{1, 2, \dots, N - k - 1\}$; for $l \in \{1, 2, \dots, k\}$ it follows that $f_{W_N}(l) \leq q^l \leq q^k \leq q^{k+1} - r = f_{W_N}(k+1)$, hence $C(W_N) = f_{W_N}(k+1) = q^{k+1} - r$. Because it is not possible for a word of length $N = q^{k+1} + k - r$, with $r \in \{1, 2, \dots, q^{k+1} - q^k\}$ to have the maximal complexity greater than $q^{k+1} - r$, it follows that $K(N) = q^{k+1} - r = N - k$.

Proposition 2. *If $\text{Card}(\mathcal{A}) = q$ and $q^k + k < N < q^{k+1} + k + 1$ then $R(N) = \{k+1\}$; if $N = q^k + k$ then $R(N) = \{k, k+1\}$.*

Proof. In the first part of the proof of Proposition 1, we proved for $N = q^{k+1} + k$, $k \geq 1$, the existence of a word W of length N for which $K(N) = f_W(k+1) = N - k$. This means that $k+1 \in R(N)$. For the word W , as well as for any other word w of length N , we have $f_w(l) < f_W(k+1)$, $l \neq k+1$, because of the special construction of W , which contains all the words of length $k+1$ in the most compact way. It follows that $R(N) = \{k+1\}$.

As in the second part of the proof of Proposition 1, we consider $N = q^{k+1} + k - r$ with $r \in \{1, 2, \dots, q^{k+1} - q^k\}$ and the word W_N for which $K(N) = f_{W_N}(k+1) = q^{k+1} - r$. We have again $k+1 \in R(N)$. For $l > k+1$, it is obvious that the complexity function of W_N , or of any other word of length N , is strictly less than $f_{W_N}(k+1)$. We examine now the possibility of finding a word W with $f_W(k+1) = N - k$ for which $f_W(l) = N - k$ for $l \leq k$. We have $f_W(l) \leq q^l \leq q^k \leq q^{k+1} - r$, hence the equality $f_W(l) = N - k = q^{k+1} - r$ holds only for $l = k$ and $r = q^{k+1} - q^k$, that is for $N = q^k + k$. We show that for $N = q^k + k$ we have indeed $R(N) = \{k, k+1\}$. If we start with Martin's word of length $q^k + k - 1$ (or with another de Bruijn word) and add to this any letter from $\mathcal{A}$, we obtain obviously a word V of length $N = q^k + k$, which contains all the q^k words of length k and $q^k = N - k$ words of length $k+1$, hence $f_V(k) = f_V(k+1) = K(N)$.

Remark 2. Having in mind the algorithm given by Martin [7] (or other more efficient algorithms), words w with maximal complexity $C(w) = K(N)$ can be easily constructed for each N and for both situations in Proposition 2.

3 De Bruijn graphs and trees

In the previous section the global maximal complexity $K(N)$ for words of length N was obtained, as well as the set of points $R(N)$ where $K(N)$ is equal to the maximal value of the subword complexity of certain words of length N . To this aim we used a special word constructed by Martin [7], which is one of the de Bruijn words. A de Bruijn word for given q and k is a word over an alphabet with q letters, containing all k -length words exactly once. The length of such a word is $q^k + k - 1$.

In order to tackle the problem of finding the number of the words for which the global maximal complexity is attained, we shall use the de Bruijn graphs and trees.

For a q -letter alphabet $\mathcal{A}$ the de Bruijn graph is defined as:

$$B(q, k) = (V(q, k), E(q, k))$$

with $V(q, k) = \mathcal{A}^k$ as the set of vertices, and $E(q, k) = \mathcal{A}^{k+1}$ as the set of directed arcs. There is an arc from $x_1x_2 \dots x_k$ to $y_1y_2 \dots y_k$ if $x_2x_3 \dots x_k = y_1y_2 \dots y_{k-1}$, and this arc is denoted by $x_1x_2 \dots x_ky_k$. See Fig. 1 and 2 for $B(2, 2)$ and $B(2, 3)$. The de Bruijn graphs $B(q, k)$ are nonplanar for $k \geq 4$, $q \geq 2$.

In the de Bruijn graph $B(q, k)$ a path (i. e. a walk with distinct vertices)

$$a_1a_2 \dots a_k, \quad a_2a_3 \dots a_{k+1}, \quad \dots, \quad a_{r-k+1}a_{r-k+2} \dots a_r \quad (r > k)$$

corresponds to an r -length word $a_1a_2 \dots a_k a_{k+1} \dots a_r$, which is obtained by adding, in turn, to the vertex $a_1a_2 \dots a_k$ the last letter of the following vertices in the path. For example in $B(2, 3)$ the path 001, 010, 101 corresponds to the word 00101. Every maximal length path in the graph $B(q, k)$ (which is a Hamiltonian one) corresponds to a de Bruijn word.




Fig. 1: The de Bruijn graph $B(2, 2)$.

In the directed graph $B(q, k)$ there always exists an Eulerian circuit because it is connected and all its vertices have the same indegree and outdegree q . An Eulerian circuit in $B(q, k)$ is a Hamiltonian path in $B(q, k + 1)$ (which always can be continued in a Hamiltonian cycle). For example in $B(2, 2)$ the following walk: 000, 001, 010, 101, 011, 111, 110, 100 represents an Eulerian circuit, which in $B(2, 3)$ is a Hamiltonian path.




Fig. 2: The de Bruijn graph $B(2, 3)$.

In order to study the number of words in $\mathcal{A}^k$ which have the maximal complexity equal to the global maximal complexity $K(N)$ we shall introduce the so-called de Bruijn trees. A *de Bruijn tree* $T(q, w)$ with the root $w \in \mathcal{A}^k$ is a q -ary tree defined recursively as follows:

- i.* The k -length word w over the alphabet $\mathcal{A} = \{e_1, e_2, \dots, e_q\}$ is the root of $T(q, w)$.
- ii.* If at any step of the recursive construction of the tree, $x_1 x_2 \dots x_k$ is a (temporary) leaf (a vertex with outdegree equal to 0), then each word among $x_2 x_3 \dots x_k e_1, x_2 x_3 \dots x_k e_2, \dots, x_2 x_3 \dots x_k e_q$ which is not in the path from the root to $x_1 x_2 \dots x_k$ will be a descendant of $x_1 x_2 \dots x_k$.
- iii.* The rule *ii* is applied as many times as it is possible.

A path is maximal if we cannot add an arc to its beginning or to its end without destroying the path property. If a maximal path is of maximal length then it is a Hamiltonian one. In any de Bruijn tree each branch is a maximal path in the de Bruijn graph $B(q, k)$ which begins with the root, and all maximal paths beginning with the root occur. For the de Bruijn trees $T(2, 000)$, $T(2, 001)$, $T(2, 010)$ and $T(2, 100)$ see Fig. 3–6. The word obtained by Martin’s algorithm corresponds to the branch of maximal length in the right side of the de Bruijn tree $T(2, 001)$.

4 Methods to compute $M(N)$

The number $M(N)$ of the words of length N for which the maximal complexity is equal to the global maximal complexity $K(N)$ can be expressed




Fig. 3: De Bruijn tree $T(2, 000)$.




Fig. 4: De Bruijn tree $T(2, 001)$.




Fig. 5: De Bruijn tree $T(2, 100)$.




Fig. 6: De Bruijn tree $T(2, 010)$.

both in terms of certain paths in a de Bruijn graph and of some vertices in the de Bruijn trees.

Proposition 3. *If $\text{Card}(\mathcal{A}) = q$ and $q^k + k \leq N \leq q^{k+1} + k$ then $M(N)$ is equal to the number of different paths of length $N - k - 1$ in the de Bruijn graph $B(q, k + 1)$.*

Proof. From Propositions 1 and 2 it follows that the number $M(N)$ of the words of length N with global maximal complexity is given by the number of words $w \in \mathcal{A}^N$ with $f_w(k + 1) = N - k$. It means that these words contain $N - k$ subwords of length $k + 1$, all of them distinct. To enumerate all of them we start successively with each word of $k + 1$ letters (hence with each vertex in $B(q, k + 1)$) and we add at each step, in turn, one of the symbols from $\mathcal{A}$ which does not duplicate a word of length $k + 1$ which has already appeared. Of course, not all of the trials will finish in a word of length N , but those which do this, are precisely paths in $B(q, k + 1)$ starting with each vertex in turn and having the length $N - k - 1$. Hence to each word of length N with $f_w(k + 1) = N - k$ we can associate a path and only one of length $N - k - 1$ starting from the vertex given by the first $k + 1$ letters of the initial word; conversely, any path of length $N - k - 1$ will provide a word w of length N which contains $N - k$ distinct subwords of length $k + 1$.

Remark 3. The number of words of length N having global maximal complexity can be also expressed by means of certain vertices in the de Bruijn trees. $M(N)$ is equal to the number of vertices at the level $N - k - 1$ in the set $\{T(q, w) \mid w \in \mathcal{A}^{k+1}\}$ of the de Bruijn trees. (The level of the root is considered to be 0, its descendants are on level 1 etc.)

The other four trees corresponding to the de Bruijn graph $B(2, 3)$ are mirror images of those in Fig. 3–6; we obtain, for example, $M(6)$ by doubling the number of vertices at level 3 in Fig. 3–6, i. e. $M(6) = 2 \cdot 18 = 36$. Similarly $M(7) = 2 \cdot 21 = 42$ is obtained by doubling the number of vertices at level 4, and so on up to $M(10) = 2 \cdot 8 = 16$ (using the vertices at level 7). These results are in accordance with those given in Table 1 obtained by counting all possible words with maximal complexity.

A formula for the number $M(N)$ of the words whose maximal complexity is equal to the global maximal complexity $K(N)$ can be given for the special case of de Bruijn words.

Proposition 4. *If $N = 2^k + k - 1$ then $M(N) = 2^{2^{k-1}}$.*

Proof. The number of distinct Hamiltonian cycles in the de Bruijn graph $B(2, k)$ is equal to $2^{2^{k-1}-k}$ [2]. With each vertex of a Hamiltonian cycle a de Bruijn word (containing all the factors of length k) begins,

which has maximal complexity, so $M(N) = 2^k \cdot 2^{2^{k-1}-k}$, which proves the proposition. (In [3] the number of circular de Bruijn words is found, which corresponds to the number of Hamiltonian cycles in de Bruijn graphs).

A generalization for $q \geq 2$ can be proved in a similar way using the results in [1].

Proposition 5. *If $N = q^k + k - 1$ then $M(N) = (q!)^{q^{k-1}}$.*

In Proposition 1, respectively Proposition 2, we have determined for each natural number N the value of the global maximal complexity $K(N)$, respectively the set of values i for which there exists a word of length N with $K(N)$ subwords of length i . To obtain a general formula for $M(N)$ for each natural number N is still an open problem.

References

- [1] T. van Aardenne-Ehrenfest, N. G. de Bruijn, *Circuits and trees in oriented linear graphs*, Simon Stevin 28 (1951), 203–217.
- [2] J. Bond, A. Iványi, *Modelling of interconnection networks using de Bruijn graphs*, Third Conference of Program Designer, Ed. A. Iványi, Budapest, 1987, 75–87.
- [3] N. G. de Bruijn, *A combinatorial problem*, Nederl. Akad. Wetensch. Proc. 49 (1946), 758–764.
- [4] M. Heinz, *Zur Teilwortkomplexität für Wörter und Folgen über einem endlichen Alphabet*, EIK 13 (1977), 27–38.
- [5] M. Lothaire, *Combinatorics on words*, Addison-Wesley, Reading, MA, 1983.
- [6] A. de Luca, *On the combinatorics of finite words*, Theor. Comput. Sci. 218 (1999), 13–39.
- [7] M. H. Martin, *A problem in arrangements*, Bull. A. M. S. 40 (1934), 859–864.